

Distribution Of Polychlorinated Biphenyls (PCBs) in Sediments From Fish Farming Cages In Different Areas Of Shatt Al-Arab Basrah /Iraq

Balqees S. Al-Ali , Dhafar Dhaher Al- Khion

Marine science center , University of Basrah , Iraq .

Abstract:

Background: Polychlorinated Biphenyls (PCBs) are found in large quantities in the environment and are poisonous, persistent, and bioaccumulative due to their high stability and propensity to build up in living things' bodies, which can have detrimental consequences on health. **Aim of study:** study the distribution of polychlorinated biphenyls (PCBs) in sediments from fish farming cages in different areas of Shatt al-Arab Basra/Iraq. **Methodology:** Composite sediment samples were collected from fish farming cages in different areas of Shatt al-Arab .Five stations were chosen along Shatt Al-Arab during (2020-2021). Sediment samples are wrapped with aluminums foil, kept in refrigerator until determination . **Results:** Current investigation, showed that high significant differences between the seasons ($P < 0.05$) among all studied group of the PCBs, in different areas of Shatt al-Arab, but this study was found that the concentrations were higher in Al-Hartha compared to other areas. **Conclusions:** The study analyzed $\sum 13$ PCBs concentrations in sediment samples from five stations in Shatt Al-Arab from 2020-2021. Results showed significant seasonal differences, with Al-Hartha having higher concentrations, indicating the influence of environmental and geographical factors on PCB levels.

Keywords : Polychlorinated Biphenyls ,PCBs, Sediments,Shatt AL-Arab River, Iraq

Introduction

Water source sediments are a secondary source of contaminants that have an impact on the local organisms. Because these creatures consume foods that contain hydrophobic chemicals, the contaminants are transferred to higher trophic levels ([Lee et al.,2001](#)). When urban and industrial garbage is discarded into rivers, lakes, and coastal waters, organic contaminants end up in the water. Compared to the ground, these chemicals are more concentrated in the surface layer. Due to their poor water solubility, they flow with rivers and floods in the direction of the sediments ([Al-Asadi and Al-Hejuje ,2019](#)). These chemical molecules have a severe poisonous effect and are extremely unstable, making them difficult to biodegrade or break down ([Al-Hejuje et al.,2017](#)). Rain, floods, and river runoff can cause a variety of hydrophobic contaminants, such as polychlorinated biphenyls, to build up in sediments ([Smith et al. ,1988](#)).

These organic molecules have a strong poisonous effect, are not particularly stable, and are difficult to break down or biodegrade ([Habeeb et al. ,2015](#)). "PCBs" or polychlorinated biphenyls are compounds with the molecular formula $C_{12}H_{10}$ or $(C_6H_5)_2$. Two benzene rings joined by a single carbon-carbon bond make up its chemical structure. One to ten chlorine atoms can be attached to the hydrogen atoms in the benzene rings, replacing the hydrogen atoms to differing degrees ([Adeyemi et al. ,2009](#)). About 130 different kinds of PCBs exist. At least fifty uniform PCB compounds may be included in commercial products and oil mixes ([You et al. ,2011](#)).

PCBs are persistent, extremely harmful compounds that are found in large quantities in the environment and bioaccumulate. These substances may be deposited into the environment through volatilization, uncontrolled leaking, leaching from soil, or industrial operations ([Combi et al. ,2016](#)). After leaking from systems that contain them, PCBs move from terrestrial sources to aquatic ecosystems through direct entry points. PCBs are chemically stable due to their lipophilic nature; they only marginally alter during the sedimentation process and accumulate by attaching themselves to particles of organic matter ([Gardes et al. ,2020](#)). Domestic and industrial wastewater are two of the many ways that PCBs used on land can seep into the aquatic environment ([Wang et al. ,2010](#)).

Recently, the amount of PCBs in water bodies has grown to be a serious issue for water quality. This substance interacts with organic matter and builds up in sediments when it enters lakes or rivers. This organic stuff is then consumed by aquatic creatures that reside in these sediments, including phytoplankton, crustaceans, and snails. As a result, fish consume these creatures and absorb PCBs. This implies that PCB contaminants may also be consumed by humans ([Iniaghe and Kpomah ,2022](#)). A high adsorption coefficient in sediments and an elevated enrichment factor in aquatic species are the effects of PCBs' restricted solubility in water ([Raghad et al.,2020](#)).

High quantities of polychlorinated biphenyls (PCBs) have been found at certain of the Tigris River's stations, according to several studies that have examined the sediments of the river. This could be the consequence of human actions that contribute to the release of PCBs into the environment, such as the use of pesticides and fertilizers on adjacent agricultural land and household garbage. This implies that these substances have been sucked up and stored in the sediments (AL-Hasani et al.,2024). Another study examined the levels of PCBs in sediment samples from the Shatt al-Arab River in Iraq. The findings showed that highly chlorinated congeners, such as tri- and hexa-PCBs, predominated over the other congeners (Al-Zabad et al.,2024). Sediments from the eastern marshes of Al-Hammar, Iraq, contained PCBs. Throughout the year, the current investigation demonstrated differences in PCB concentrations between sites and seasons. Sediments can also be regarded as an indicator of the degree of pollution, even if water analysis is a useful method for evaluating river contamination. Pollutants are stored in the surface sediments of the rivers under study, and as environmental conditions change, the buildup of these pollutants in the sediments could contaminate the water (Al-Hejuje et al.,2015).

Studying PCBs has become crucial in the field of environmental research because of their strong chemical affinity for lipids, their possible carcinogenic effects, and the bioaccumulation that results (Cortazar et al.,2002). This present study aims to examine the distribution of polychlorinated biphenyls (PCBs) in the sediments of fish farming cages in different parts of Shatt al-Arab in Basra, Iraq.

Methodology

Sediment sampling

Composite sediment samples were collected from fish farming cages in different areas of Shatt al-Arab. Five stations were chosen along Shatt Al-Arab during (2020-2021). (Al-musahab, Abu Al-Khasib, Alharitha , Al-Nashwa and Al-Nujaira) (Fig1). Sediment samples are wrapped with aluminums foil, kept in refrigerator until determination

Figure 1: Map of the study area.

Sample extraction, clean-up and determination

A measured weight of sample (approximately 2 to 30 g of sediment) is extracted using METHOD 8082A (2007). Sediment sample are extracted with Hexane-acetone (1:1), using (Soxhlet), cleanup with Florisil, after cleanup, the extract is analyzed by injecting a measured aliquot into a GC-mass (Agilent5975 , USA) , Type of column: 30-m x 0.25-mm ID fused-silica capillary column ,HP-5MS , 0.25 μ m film thickness.

Statistical Analysis

Data of current study were analyzed by using F test (ANOVA) used to compare between three numeric variables or more. Numeric data were described by mean $\pm$ standard deviation (SD). A level of significance of $\alpha=0.05$ was applied to test. (SPSS v.26 and excel 2013) programs used to analyze current data..

Results

The current study, showed that the concentration of PCB compounds in different areas of Shatt al-Arab. Five stations were chosen along Shatt Al-Arab during (2020-2021). (Al-musahab, Abu Al-Khasib, Alharitha , Al-Nashwa and Al-Nujaira) during Autumn, winter, and winter.

In Present study, ,during Autumn , the highest concentration of PCB 18 (1.85 ng/g d.w.) occurred at Abu Al-Khasib while, the concentrations were low in the other PCP types across most stations in this study,tables (1) and figures (2).

Table (1): PCB concentrations (ng/g dry weight) in the sediments of the study stations during Autumn.

Figure 2: Concentration and types of PCBs in different areas of Shatt al-Arab during Autumn.

In winter, the highest concentration of PCB 44 (0.828 ng/g d.w.) occurred at Nujayrat Dakhil while the concentrations were low in the other PCP types across most stations in this study,tables (2) and figures (3).

Table (2): PCB concentrations (ng/g dry weight) in the sediments of the study stations during winter**Figure 3:** Concentration and types of PCBs in different areas of Shatt al-Arab during Winter.

In Spring, the highest concentration of PCB 101(4.257 ng/g d.w.) occurred at alharitha, while,PCB 28 and PCB 149 were lowest in all stations in this study,tables (3) and figures (4).

Table (3): PCB concentrations (ng/g dry weight) in the sediments of the study stations during Spring**Figure4 :** Concentration and types of PCBs in different areas of Shatt al-Arab during Spring.

In Summer, the highest concentration of PCB 52 (3.35 ng/g d.w.) occurred at alharitha, while,PCB 28 and PCB 149 were lowest in all stations in this study,table (4) and figures (5).

Table (4): PCB concentrations (ng/g dry weight) in the sediments of the study stations during Summer**Figure 5:** Concentration and types of PCBs in different areas of Shatt al-Arab during Summer.

In Present investigation,the statistical analysis using one-way ANOVA showed that high significant differences between the seasons ($P<0.05$) among all studied group of the PCBs, in different areas of Shatt al-Arab, but this study was found that the concentrations were higher in Al-Hartha compared to other areas , table(5) and figure(6).

Table(5) : Mean concentrations of PCBs in different areas of Shatt al-Arab during the four seasons**Figure 6:** Mean concentrations of PCBs in different areas of Shatt al-Arab during the four seasons.

Discussion

The properties of PCBs compounds, including heat resistance, low flammability, and high dielectric constant, make them suitable for multiple industrial applications, such as refrigerants and insulation fluids for condensers and electrical

transformers. It is also used in the manufacture of PVC wrappers, pesticides, cutting oils, flame retardants, hydraulic fluids, sealants, adhesives, paints, and carbon-free printing paper (BRINKMAN and de KOK, 1980). These sources may supply rivers with high concentrations of PCBs compounds. PCBs used on land can travel to the aquatic environment through various pathway routes including wastewater discharged by industry, municipal wastewater, river input, air transport and sedimentation (Yang et al., 2012), once transported to the aquatic environment, PCBs can be absorbed by molecules and then incorporated into sediment because they were difficult to dissolve in water, which can lead to pollution (Yang et al., 2011), sediments are a clear evidence of aquatic environmental pollution as large quantities of PCBs show that sediment is a reservoir and will be a clear environmental deposit of these pollutants in the aquatic environment.

Polychlorinated Biphenyls (PCB) are persistent, toxic, bioaccumulative, and widely distributed in the environment because of their high stability and tendency to accumulate in the bodies of living organisms, causing serious health effects (AL-Hasani et al., 2024).

The results showed that the highest concentration of PCBs was recorded during spring, followed by summer, then autumn, while the lowest concentration was detected in winter. Additionally, it was found that the concentrations were higher in Al-Hartha compared to other areas, including Abu Al-Khasib, Al-Nashwa, Najiran, and Al-Mishkhab. This increase may be attributed to the fact the sediment texture in Al-Hartha station was silty clay, which is different from the rest of the stations that were sand-silt sediments, as the silt-clay sediments accumulated organic compounds; in addition, the smallest size of the grains in the sediments can accumulate greater amounts of PCBs. In addition, the values of PCBs in clay sediments have increased more than those in sandy sediments (Al-Zabad et al., 2021). A study by Barakat et al. (2013) indicated that sediments that are close to cities often have high concentrations of PCBs according to the local use of these pollutants. The results showed that the increase in the concentrations of PCBs in the sediments of Alharitha station in the summer may be due to the main roads adjacent to Shatt al-Arab, where they are affected by waste from means of transportation and activated fishing boats that increase water pollution and then settle in sediments.

This finding was consistent with the study conducted by AL-Hasani et al. (2024), where PCB levels were found to increase during the summer season. These seasonal and geographical variations can be explained based on environmental, hydrodynamic, and anthropogenic factors affecting the deposition and distribution of PCBs in the Shatt Al-Arab environment, particularly in fish farming cages. The Effect of Temperature on PCB Deposition and Accumulation: PCBs are hydrophobic compounds, meaning they tend to bind to organic matter and sediments rather than remain dissolved in water. As temperature increased during spring and summer, the rates of chemical and biological reactions affecting the movement and redistribution of these compounds were also elevated. Several key factors may explain this increase: Enhanced microbial activity: Higher temperatures led to an increase in bacterial activity, which may have contributed to the decomposition of organic matter in sediments, leading to the release and redistribution of PCBs (Jazaa, 2009).

Volatilization of lighter PCB congeners: Some PCBs evaporated or migrated from water to sediments at higher rates due to increased temperature. Increased biological activity of fish and other aquatic organisms, which resulted in sediment disturbance and redistribution of pollutants such as PCBs.

The Effect of Water Flow and River Currents: During winter, the water flow rates and river currents in Shatt Al-Arab were higher due to increased rainfall and rising water levels. This increase led to: Dilution of PCB concentrations in sediments, as stronger currents dispersed contaminants. The transport of some contaminated sediments away from the sampling sites, reducing the accumulation of PCBs in those areas.

In contrast, during spring and summer, water flow decreased, and evaporation rates were higher, leading to increased PCB concentrations in sediments due to reduced dilution and greater retention of pollutants.

The Effect of Geographical Factors on PCB Concentration Differences Between Areas: The results indicated that the highest PCB concentrations were found in Al-Hartha, compared to other areas such as Abu Al-Khasib, Al-Nashwa, Najiran, and Al-Mishkhab. This variation can be attributed to several factors: Proximity to industrial pollution sources (Mills and Chichester, 2005) : Al-Hartha is an industrial area containing factories and oil refineries, which may have contributed to the discharge of large amounts of pollutants, including PCBs, into Shatt Al-Arab.

Lower water flow compared to other regions: Reduced water currents in some parts of Al-Hartha led to higher PCB accumulation in sediments due to lower dispersion and dilution rates.

Direct impact of fish farming cages: Waste from fish and feed materials may have promoted the accumulation of organic matter, enhancing PCB absorption in sediments (Al-Asadi and Al-Hejuje, 2019).

The Effect of Human and Industrial Activities: During spring and summer, industrial and agricultural activities surrounding Shatt Al-Arab increased, leading to greater discharge of untreated industrial wastewater and agricultural runoff containing organic pollutants such as PCBs.

Summer was also characterized by intensified fishing and fish farming activities, which may have caused sediment disturbances and redistribution of pollutants. In Al-Hartha, industrial activity was likely more significant, explaining the higher PCB levels in sediments compared to other locations (Hsu, 2004).

Conclusion : The study determined and analyzed the concentrations of $\Sigma 13$ PCBs compound in sediment samples at five stations were chosen along Shatt Al-Arab during (2020-2021). Current investigation, showed that high significant differences between the seasons ($P < 0.05$) among all studied group of the PCBs, in different areas of Shatt al-Arab, but this study was found that the concentrations were higher in Al-Hartha compared to other areas. These findings reflect the strong influence of environmental and geographical factors, such as temperature, water flow, and industrial and agricultural activities, on PCB levels in sediments.

References

- Adeyemi, D., Ukpo, G., Anyakora, C., & Uyimadu, J. (2009). Polychlorinated biphenyl in fish samples from Lagos Lagoon, Nigeria. *African Journal of Biotechnology*, 8(12). <http://www.academicjournals.org/AJB>
- Al-Asadi, A.A. & Al-Hejuje, M.M. (2019). Application of organic pollution index (OPI) to assess the water quality of Al-Chibayish Marsh, Southern Iraq. *Marsh Bull.*, 14(1): 57-64.
- AL-Hasani ,AL-Hasanayn M.; Al-Hejuje, Makia M.& Al-saad ,H. T. (2024). Polychlorinated Biphenyls (PCB) in sediments of the southern part of Tigris River-Iraq. *Marsh Bulletin* , Vol.19 No.1 65-73.
- Al-Hejuje M.M.; Al-Saad H.T.; and Hussain, N.A.(2015). Total petroleum hydrocarbons (TPHs), n-alkanes, and polynuclear Aromatic Hydrocarbons (PAHs) in Sediments of Shatt Al-Arab River, Part 2. *Global Journal of Biology, Agriculture & Health Sciences (G.J.B.A.H.S.)*, 4(1):95-100. <https://www.scholarscentral.com/>
- Al-Hejuje, M. M., Al-saad, H. T., & Hussain, N. A. (2017). Assessing the Organic Pollution and Aquaculture Activity of Surface Water at Shatt-Al-Arab Estuary Southern of Iraq. *Indonesian Journal of Marine Sciences/Ilmu Kelautan*, 22(4).
- Al-Zabad, R. A., Al-Khafaji, A. H., & Al- Saad, H. T. (2021). Concentrations Of Polychlorinated Biphenyls (PCBs) In *Phragmites australis* of Shatt AL-Arab River, Basrah Governorate South of Iraq. *Al- Qadisiyah Journal of Pure Science*, 26(4), 199–209. <https://doi.org/10.29350/qjps.2021.26.4.142.2>
- Al-Zabad, R. A. Al-Khafaji, A. H. and AL-Saad H. T. (2024). Analyzing and distribution of polychlorinated biphenyls (PCBs) in sediments along Shatt Al-Arab estuary ,Iraq. *Ecology and environmental technology* No.4/3(78)p.33-38.
- Barakat, A. O., Khairy, M., & Aukaily, I. (2013). Persistent organochlorine pesticide and PCB residues in surface sediments of Lake Qarun, a protected area of Egypt. *Chemosphere*, 90(9), 2467–2476. <https://doi.org/10.1016/j.chemosphere.2012.11.012>
- Brinkman, U. A., and De Kok, A. (1980). Halogenated Biphenyls, Terphenyls, Naphthalenes, Dibenzodioxins and Related Products. *Chemosphere*, 21(8), 1007–1016.
- Combi, T., Miserocchi, S., Langone, L., & Guerra, R. (2016). Polychlorinated biphenyls (PCBs) in sediments from the western Adriatic Sea: Sources, historical trends, and inventories. *Science of The Total Environment*, 562, 580–587. <https://doi.org/10.1016/j.scitotenv.2016.04.086>
- Cortazar, E., Zuloaga, O., Sanz, J., Raposo, J. C., Etzebarria, N., & Fernandez, L. A. (2002). Multisimplex optimization of solid-phase microextraction–gas chromatographic–mass spectrometric determination of polycyclic aromatic hydrocarbons, polychlorinated biphenyls, and phthalates from water samples. *Journal of Chromatography A*, 978(1-2), 165-175.
- Gardes, T., Portet-Koltalo, F., Debret, M., Humbert, K., Levailant, R., Simon, M., & Copard, Y. (2020). Temporal trends, sources, and relationships between sediment characteristics and polycyclic aromatic hydrocarbons (PAHs) and polychlorinated biphenyls (PCBs) in sediment cores from the major Seine estuary tributary, France. *Applied Geochemistry*, 122, 104749. <https://doi.org/10.1016/j.apgeochem.2020.104749>
- Habeeb, M. A., Al-Bermani, A. K., & Salman, J. M. (2015). Environmental study of water quality and some heavy metals in water, sediment and aquatic macrophytas in lotic ecosystem, Iraq. *Mesopotamia Environmental Journal*, 1(2), 66–84.

- Hsu, S. (2004). Removal Polycyclic Aromatic Hydrocarbons (phenanthrene & pyrene) in aquatic phase by *Ceratophyllum demorsum* & *Naja gramunea*. Master thesis. Marine Environment & Engineering. chinese.
- Iniaghe, P. O., & Kpomah, E. D. (2022). Polychlorinated biphenyls (PCBs) in water and sediments from the Udu River, Niger Delta, Nigeria: concentration, distribution and risk assessment. *Journal of Environmental Exposure Assessment*, 1(3), 20. <https://doi.org/10.20517/jeea.2022.19>
- Jazaa, S. H. (2009). A study of physical, chemical and bacteriological properties of water of Al-Kahlaa river in maysan governorate/Iraq. M. Sc. Thesis. College of Science. University of Basrah. pp: 67
- Lee, K. T., Tanabe, S., & Koh, C. H. (2001). Contamination of polychlorinated biphenyls (PCBs) in sediments from Kyeonggi Bay and nearby areas, Korea. *Marine Pollution Bulletin*, 42(4), 273-279.
- Mills, L. J., & Chichester, C. (2005). Review of evidence: are endocrine-disrupting chemicals in the aquatic environment impacting fish populations? *Science of the Total Environment*, 343(1-3), 1-34.
- Raghad, S. J., Ahmed Y., Zuhair A. A., Abdul Zahra A. and Al-Saad, H. T. (2020). An integrative study to determine the Bioavailability of heavy metals in the soil, Basrah, Iraq. *Eco. Env. & Cons.* 25 (4) : 2019; pp. (35-42) .
- Smith, J. A., Witkowski, P. J., & Fusillo, T. V. (1988). Manmade organic compounds in the surface waters of the United States: a review of current understanding.
- Yang, H., Xue, B., Jin, L., Zhou, S., & Liu, W. (2011). Polychlorinated biphenyls in surface sediments of Yueqing Bay, Xiangshan Bay, and Sanmen Bay in East China Sea. *Chemosphere*, 83(2), 137-143. <https://doi.org/10.1016/j.chemosphere.2010.12.070>
- Yang, H., Zhuo, S., Xue, B., Zhang, C., & Liu, W. (2012). Distribution, historical trends and inventories of polychlorinated biphenyls in sediments from Yangtze River Estuary and adjacent East China Sea. *Environmental Pollution*, 169, 20-26. <https://doi.org/10.1016/j.envpol.2012.05.003>.
- You, H., Ding, J., Zhao, X.-S., Li, Y.-F., Liu, L.-Y., Ma, W.-L., Qi, H., & Shen, J.-M. (2011). Spatial and seasonal variation of polychlorinated biphenyls in Songhua River, China. *Environmental Geochemistry and Health*, 33(3), 291-299. <https://doi.org/10.1007/s10653-010-9341-7>
- Wang, W., Bai, J., Zhang, G., Jia, J., Wang, X., Liu, X., & Cui, B. (2019). Occurrence, sources and ecotoxicological risks of polychlorinated biphenyls (PCBs) in sediment cores from urban, rural and reclamation-affected rivers of the Pearl River Delta, China. *Chemosphere*, 218, 359-367. <https://doi.org/10.1016/j.chemosphere.2018.11.046>